

Signal for CP violation in $B^\pm \rightarrow P\bar{P}\pi^\pm$ decays

B. Bajc^a, S. Fajfer^a, R. J. Oakes^b, T.N. Pham^c and S. Prelovšek^a

^aJ. Stefan Institute, Jamova 39, P. O. Box 3000, 1001 Ljubljana, Slovenia

^bDepartment of Physics and Astronomy, Northwestern University, Evanston, IL 60208, U.S.A.

^cCentre de Physique Theorique, Centre National de la Recherche Scientifique, UPR A0014, Ecole Polytechnique, 91128 Palaiseau Cedex, France

We analyze the partial rate asymmetry in $B^\pm \rightarrow P\bar{P}\pi^\pm$ decays ($P = \pi^+, K^+, \pi^0, \eta$) which results from the interference of the nonresonant decay amplitude and the resonant amplitude for $B^\pm \rightarrow \chi_{c0}\pi^\pm$ followed by the decay $\chi_{c0} \rightarrow P\bar{P}$. The CP violating phase γ can be extracted from the measured asymmetry. We find that the partial rate asymmetry for $B^\pm \rightarrow \pi^+\pi^-\pi^\pm$ is $0.33 \sin\gamma$, while for $B^\pm \rightarrow K^+K^-\pi^\pm$ it amounts $0.45 \sin\gamma$.

The measurement of CP asymmetries in charged B meson decays might provide us with a first demonstration of CP violation outside the K system [1,2]. One possibility to measure the CP odd phase $\gamma = \arg(V_{ub}^*)$ has been suggested in [3]. In this work the asymmetry appears as a result of the interference of the nonresonant decay $B^\pm \rightarrow P\bar{P}\pi^\pm$ amplitude and the resonant $B^\pm \rightarrow \chi_{c0}\pi^\pm \rightarrow P\bar{P}\pi^\pm$, where $\chi_{c0}(3.4)$ is a $c\bar{c}$ scalar [3,4]. The absorptive phase necessary to observe CP violation in partial rate asymmetries is provided by the χ_{c0} width.

On the experimental side the CLEO collaboration has reported upper limits on $B^+ \rightarrow h^+h^+h^-$ nonresonant decays [5]. They found that the upper limit on the branching ratio $BR(B^+ \rightarrow \pi^+\pi^-\pi^+) \leq 4.1 \times 10^{-5}$ and $BR(B^+ \rightarrow K^+K^-\pi^+) \leq 7.5 \times 10^{-5}$. The authors of [3] estimated theoretically that the branching ratio for $B^+ \rightarrow \pi^+\pi^-\pi^+$ ranges from 1.5×10^{-5} to 8.4×10^{-5} .

Motivated by the theoretical expectation and the experimental result, we investigate the $B^\pm \rightarrow P\bar{P}\pi^\pm$ nonresonant decay widths, where $P = \pi^+, K^+, \pi^0, \eta$. We also study the resonant decay amplitudes of $B^\pm \rightarrow P\bar{P}\pi^\pm$ arising from χ_{c0} decays into $P\bar{P}$, $P = \pi^+, K^+, \pi^0, \eta$. The branching ratios of the χ_{c0} decays into these modes have been reported [6]. We point out that we are interested only in the kinematical

region where the $P\bar{P}$ invariant mass is close to the χ_{c0} mass, like in [3]. Thus $P\bar{P}$ arising from resonances such as ρ , and other possible resonances will not be considered in this paper. In the analysis of the nonresonant decay amplitude of $B^\pm \rightarrow P\bar{P}\pi^\pm$ we use the factorization approximation, which shows that the main contribution comes from the product $\langle P\bar{P} | (\bar{u}b)_{V-A} | B^- \rangle \langle \pi^- | (\bar{d}u)_{V-A} | 0 \rangle$ ($(\bar{q}_1 q_2)_{V-A}$ denotes $\bar{q}_1 \gamma_\mu (1 - \gamma_5) q_2$). For the calculation of the matrix element $\langle P\bar{P} | (\bar{u}b)_{V-A} | B^- \rangle$ we use the results obtained in [7], where the nonresonant $D^+ \rightarrow K^-\pi^+\nu$ decay was analyzed. The experimental result for the branching ratio of the nonresonant $D^+ \rightarrow K^-\pi^+\nu$ decay was successfully reproduced within a framework (hybrid model) which combines the heavy quark effective theory (HQET) and the chiral Lagrangian (CHPT). The combination of heavy quark symmetry and chiral symmetry was quite successful in the analysis of D meson semileptonic decays [8,9], and references therein. The heavy quark symmetry is expected to work better for B mesons [8]. However, new difficulties might appear in B decays, due to the large energies of light mesons in the final state. It is known, however, that the combination of HQET and CHPT is valid at small recoil momentum. The details of our approach developed for D meson semileptonic decays, can be found in [7,9]. We systematically apply the hybrid model in the

calculation of the $B^\pm \rightarrow P\bar{P}\pi^\pm$ amplitude and we find that in a weak transition of $B \rightarrow P\bar{P}$ new important contributions arise, which were not taken into account in [3].

The weak effective Lagrangian for the non-leptonic Cabibbo suppressed B meson decays is given by [3]

$$\mathcal{L}_w = -\frac{G_F}{\sqrt{2}} V_{ud}^* V_{ub} (a_1^{eff} O_1 + a_2^{eff} O_2) \quad (1)$$

where $a_1^{eff} \simeq 1.08$ and $a_2^{eff} \simeq 0.21$ [2], $O_1 = (\bar{u}b)_{V-A}(\bar{d}u)_{V-A}$ and $O_2 = (\bar{u}u)_{V-A}(\bar{d}b)_{V-A}$. The quark current required in the weak Hamiltonian (1) is expressed in terms of meson fields [7]. In our calculations we will follow the approach described in [7] and [9]: we use the Feynman rules for the vertices near and outside the zero-recoil region, *but we include the complete propagators instead of using the HQET propagator* [8].

Using the factorization approach [3], we analyze all possible contributions to the $B^\pm \rightarrow \pi^\pm P\bar{P}$ nonresonant amplitude. We notice that the main contribution to the amplitude for nonresonant $B^\pm \rightarrow P\bar{P}\pi^\pm$ decay comes from the matrix element of $\langle P(p_1)\bar{P}(p_2) | (\bar{u}b)_{V-A} | B^-(p_B) \rangle$. Following the analysis described in [7], we write

$$\begin{aligned} \langle \pi^-(p_1)\pi^+(p_2) | \bar{u}\gamma_\mu(1-\gamma_5)b | B^-(p_B) \rangle = \\ i r(p_B - p_2 - p_1)_\mu + i w_+(p_2 + p_1)_\mu + \\ i w_-(p_2 - p_1)_\mu + 2h\epsilon_{\mu\alpha\beta\gamma} p_B^\alpha p_2^\beta p_1^\gamma. \end{aligned} \quad (2)$$

In our case only the nonresonant form factors w_-^{nr} and w_+^{nr} contribute. The contribution proportional to r is of order m_π^2 and therefore we can safely neglect it. In the case of $B^- \rightarrow \pi^-\pi^-\pi^+$ decay the form factors w_+ and w_- are given by

$$\begin{aligned} w_+^{nr}(s, t) = -\frac{g}{f_\pi^2} \frac{f_{B^*} m_{B^*}^{3/2} m_B^{1/2}}{s - m_{B^*}^2} \times \\ \left\{ 1 - \frac{1}{2m_{B^*}^2} (m_B^2 - m_\pi^2 - s) \right\} + \frac{f_B}{2f_\pi^2} - \\ \frac{\sqrt{m_B} \alpha_2}{2f_\pi^2} \frac{1}{m_B^2} (2s + t - m_B^2 - 3m_\pi^2), \end{aligned} \quad (3)$$

$$\begin{aligned} w_-^{nr}(s) = \frac{g}{f_\pi^2} \frac{f_{B^*} m_{B^*}^{3/2} m_B^{1/2}}{s - m_{B^*}^2} \times \\ \left\{ 1 + \frac{1}{2m_{B^*}^2} (m_B^2 - m_\pi^2 - s) \right\} + \frac{\sqrt{m_B} \alpha_1}{f_\pi^2}. \end{aligned} \quad (4)$$

In the above expressions $f_{B, (B^*)}$ are B meson decay constants, the parameters $\alpha_{1,2}$ are fixed in [9] using the data from semileptonic $D^0 \rightarrow \bar{K}^{*0} l \nu_l$ decay. The s , t and u are defined as usual: $s = (p_B - p_3)^2 = (p_2 + p_1)^2$, $t = (p_B - p_1)^2 = (p_1 + p_2)^2$ and $u = (p_B - p_2)^2 = (p_1 + p_3)^2$. The decay constants are related by the heavy quark symmetry [8]. We use $f_D \simeq 200 \text{ MeV}$ [9], giving $f_B \simeq 120 \text{ MeV}$. Then, we use $|V_{ub}| = 0.003$ [6]. The parameters α_1 and α_2 must be specified for B meson decays. It is easy to find that the soft scaling [8] of the axial form factors A_1 and A_2 , leads to the following relations: $\alpha_1^{DK^*} = \alpha_1^{B\rho}$, while $\alpha_2^{DK^*}/m_D = \alpha_2^{B\rho}/m_B$. We take the parameters $\alpha_{1,2}^{DK^*}$ from D meson semileptonic decays [7,9]. branching ratio $B^- \rightarrow \pi^-\pi^+\pi^-$

The parameter $g = 0.3 \pm 0.1$ is determined from $D^* \rightarrow D\pi$ decays [10]. From $D^0 \rightarrow K^- l^+ \nu$ we have found $g = 0.15 \pm 0.08$ [11]. In the present calculations we take the range $0.2 \leq g \leq 0.23$ and we select $\alpha_1^{DK^*} = -0.13 \text{ GeV}^{1/2}$, $\alpha_2^{DK^*} = -0.13 \text{ GeV}^{1/2}$, giving $3.4 \times 10^{-5} \leq BR(B^- \rightarrow \pi^-\pi^+\pi^-) \leq 3.8 \times 10^{-5}$. All other combinations of parameters give the branching ratio too large in comparison with the experimental limit. Using the same set of parameters, we find $1.4 \times 10^{-5} \leq BR(B^- \rightarrow K^- K^+ \pi^-) \leq 1.5 \times 10^{-5}$. We calculate the following limits for the branching ratios $4.6 \times 10^{-6} \leq BR(B^- \rightarrow \pi^-\pi^0\pi^0) \leq 6.1 \times 10^{-6}$ and $6.4 \times 10^{-7} \leq BR(B^- \rightarrow \pi^-\eta\eta) \leq 8.5 \times 10^{-7}$. We find that the contributions proportional to the $\alpha_{1,2}$ in the calculated branching ratio are very important.

The resonant decay amplitude for $B^- \rightarrow \chi_{0c}\pi^- \rightarrow \pi^+\pi^-\pi^-$ is determined by the narrow width approximation

$$\begin{aligned} \mathcal{M}_r(B^- \rightarrow \chi_{0c}\pi^- \rightarrow \pi^+\pi^-\pi^-) = \\ M(B^- \rightarrow \chi_{0c}\pi^-) \frac{1}{s - m_{\chi_{0c}}^2 + i\Gamma_{\chi_{0c}} m_{\chi_{0c}}} \times \\ M(\chi_{0c} \rightarrow \pi^+\pi^-) + (s \leftrightarrow t). \end{aligned} \quad (5)$$

We have used the estimation $BR(B^\pm \rightarrow \chi_{c0}\pi^\pm)/BR(\chi_{c0} \rightarrow \pi^+\pi^-) = 5 \times 10^{-7}$ found in [4]. Using the χ_{c0} data [6], we fix the remaining parameters.

Following the analysis of [3], we investigate the partial rate asymmetry in $B^\pm \rightarrow \pi^\pm P\bar{P}$. We are interested in the kinematical region, where the $P\bar{P}$ invariant mass is close to the mass of the χ_{c0} meson $m_{\chi_{c0}} = 3.415 \text{ GeV}$. The partial decay width Γ_p for $B^- \rightarrow \pi^+\pi^-\pi^-$, which contains nonresonant and resonant contributions, is obtained by the phase space integration from $s_l = (m_{\chi_{c0}} - 2\Gamma_{\chi_{c0}})^2$ to $s_u = (m_{\chi_{c0}} + 2\Gamma_{\chi_{c0}})^2$, where $\Gamma_{\chi_{c0}} = 0.014 \pm 0.005 \text{ GeV}$ is the width of χ_{c0}

$$\Gamma_p = C \int_{s_l}^{s_u} ds \int_{t_l(s)}^{t_u(s)} dt |\mathcal{M}_{nr} + \mathcal{M}_r|^2, \quad (6)$$

where $C = 1/[(2\pi m_B)^3 32]$. Similarly, $\Gamma_{\bar{p}}$ is the partial decay width for $B^+ \rightarrow \pi^+\pi^-\pi^+$ containing the nonresonant and resonant contribution. The CP-violating asymmetry is determined by

$$|A| = \left| \frac{\Gamma_p - \Gamma_{\bar{p}}}{\Gamma_p + \Gamma_{\bar{p}}} \right|. \quad (7)$$

In our calculations of the branching ratio the CHPT is used beyond its region of applicability and the obtained results should be taken with care. In the calculation of the partial decay width, however, the kinematical region is constrained to the region around the χ_{c0} resonance and therefore our calculations for the partial decay asymmetry are more reliable.

For the selected set of parameters g , $\alpha_{1,2}$ we obtain

$$A(B^+ \rightarrow \pi^+\pi^-\pi^+) = 0.33 \sin\gamma, \quad (8)$$

$$A(B^+ \rightarrow K^+K^-\pi^+) = 0.45 \sin\gamma, \quad (9)$$

while the asymmetry in the case of $B^+ \rightarrow \pi^0\pi^0\pi^+$ decay ranges from $(0.14 - 0.16) \sin\gamma$ and in the case of $B^+ \rightarrow \eta\eta\pi^+$ decay ranges from $(0.10 - 0.17) \sin\gamma$.

We have analyzed the partial rate asymmetry in $B^\pm \rightarrow P\bar{P}\pi^\pm$ decays ($P = \pi^+, K^+, \pi^0, \eta$),

which signals CP violation. The nonzero asymmetry results from the interference of the nonresonant decay amplitude and the resonant decay amplitude $B^\pm \rightarrow \chi_{c0}\pi^\pm$ followed by $\chi_{c0} \rightarrow P\bar{P}$. The asymmetry is found to be rather dependent on the choice of parameters and is $0.33 \sin\gamma$ ($0.45 \sin\gamma$) for $B^+\pi^+\pi^-\pi^+$ ($B^+ \rightarrow \pi^+K^-K^+$), while it is smaller for $B^+ \rightarrow \pi^0\pi^0\pi^0$ and $B^+ \rightarrow \eta\eta\pi^+$ decays. The calculated partial rate asymmetries give useful guidelines for the experimental searches of the size of CP - violating angle γ .

REFERENCES

1. Y. Nir, talk given at 18th International Symposium on Lepton Photon Interactions, Hamburg, Germany, July 28 - August 1 1997, hep-ph/9709301.
2. A. Ali, G. Kramer and Dai-Dian Lü, hep-ph/9805403.
3. N. G. Deshpande, G. Eilam, X. G. He and J. Trampetić, Phys. Rev. D 52 (1995) 5354.
4. G. Eilam, M. Gronau and R. R. Mendel, Phys. Rev. Lett. 74 (1995) 4984.
5. T. Bergfeld et al., Phys. Rev. Lett. 77 (1996) 4503.
6. R. M. Barnett et al., Phys. Rev. D 54 (1996) 1.
7. B. Bajc, S. Fajfer, R. J. Oakes and T. N. Pham, hep-ph/9710422, to appear in Phys. Rev. D.
8. R. Casalbuoni, A. Deandrea, N. Di Bartolomeo, R. Gatto, F. Feruglio and G. Nardulli, Phys. Rep. 281 (1997) 371.
9. B. Bajc, S. Fajfer and R. J. Oakes, Phys. Rev. D 53 (1996) 4957.
10. P. Colangelo, Proceedings of this conference.
11. B. Bajc, S. Fajfer, R.J. Oakes, S. Prelovsek, Phys. Rev. D 56 (1997) 7207.